

-1-

METHOD OF USING A MATRIX METALLAPROTEINASE INHIBITOR AND
RADIATION THERAPY AS COMBINATION THERAPY IN THE
TREATMENT OF NEOPLASIA

5 **Field of the Invention**

The present invention relates to a combination of radiation therapy and a matrix metalloproteinase (MMP) inhibitor for treatment of neoplasia disorders. More specifically, this invention relates to the use of MMP
10 inhibitors in combination with radiation therapy for treating cancer.

Background of the Invention

A neoplasm, or tumor, is an abnormal, unregulated,
15 and disorganized proliferation of cell growth. A neoplasm is malignant, or cancerous, if it has properties of destructive growth, invasiveness and metastasis. Invasiveness refers to the local spread of a neoplasm by infiltration or destruction of surrounding
20 tissue, typically breaking through the basal laminas that define the boundaries of the tissues, thereby often entering the body's circulatory system. Metastasis typically refers to the dissemination of tumor cells by lymphatics or blood vessels. Metastasis also refers to
25 the migration of tumor cells by direct extension through serous cavities, or subarachnoid or other spaces. Through the process of metastasis, tumor cell migration to other areas of the body establishes neoplasms in areas away from the site of initial appearance.

30 Cancer is now the second leading cause of death in the United States and over 8,000,000 persons in the United States have been diagnosed with cancer. In 1995,

-2-

cancer accounted for 23.3% of all deaths in the United States.

Cancer is not fully understood on the molecular level. It is known that exposure of a cell to a

5 carcinogen such as certain viruses, certain chemicals, or radiation, leads to DNA alteration that inactivates a "suppressive" gene or activates an "oncogene".

Suppressive genes are growth regulatory genes, which upon mutation, can no longer control cell growth.

10 Oncogenes are initially normal genes (called proto-oncogenes) that by mutation or altered context of expression become transforming genes. The products of transforming genes cause inappropriate cell growth. More than twenty different normal cellular genes can become
15 oncogenes by genetic alteration. Transformed cells differ from normal cells in many ways, including cell morphology, cell-to-cell interactions, membrane content, cytoskeletal structure, protein secretion, gene expression and mortality.

20 Cancer is now primarily treated with one or a combination of three types of therapies: surgery, radiation, and chemotherapy. Surgery involves the bulk removal of diseased tissue. While surgery is sometimes effective in removing tumors located at certain sites,
25 for example, in the breast, colon, and skin, it cannot be used in the treatment of tumors located in other areas, inaccessible to surgeons, nor in the treatment of disseminated neoplastic conditions such as leukemia.

Chemotherapy involves the disruption of cell
30 replication or cell metabolism. It is used most often in the treatment of breast, lung, and testicular cancer.

-3-

The adverse effects of systemic chemotherapy used in the treatment of neoplastic disease is most feared by patients undergoing treatment for cancer. Of these adverse effects nausea and vomiting are the most common and severe side effects. Other adverse side effects include cytopenia, infection, cachexia, mucositis in patients receiving high doses of chemotherapy with bone marrow rescue or radiation therapy; alopecia (hair loss); cutaneous complications such as pruritis, urticaria, and angioedema; neurological complications; pulmonary and cardiac complications in patients receiving radiation or chemotherapy; and reproductive and endocrine complications (M. Abeloff, et al., Alopecia and Cutaneous Complications, in Clinical Oncology 755-56 (Abeloff, ed. 1992).

Chemotherapy-induced side effects significantly impact the quality of life of the patient and may dramatically influence patient compliance with treatment.

Additionally, adverse side effects associated with chemotherapeutic agents are generally the major dose-limiting toxicity (DLT) in the administration of these drugs. For example, mucositis, is one of the major dose limiting toxicity for several anticancer agents, including the antimetabolite cytotoxic agents 5-FU, methotrexate, and antitumor antibiotics, such as doxorubicin. Many of these chemotherapy-induced side effects if severe, may lead to hospitalization, or require treatment with analgesics for the treatment of pain.

In general, radiation therapy is employed as potentially curative therapy for patients who present

-4-

with clinically localized disease and are expected to live at least 10 years.

For example, approximately 70% of newly diagnosed prostate cancer patients fall into this category.

5 Approximately 10% of these patients (7% of total patients) undergo radiation therapy. Approximately 80% of patients who have undergone radiation as their primary therapy have disease persistence or develop recurrence or metastasis within five years after
10 treatment. Currently, most of these radiotherapy patients generally do not receive any immediate follow-up therapy. Rather, they are monitored frequently, such as for elevated Prostate Specific Antigen ("PSA"), which is the primary indicator of recurrence or metastasis in
15 prostate cancer.

The adverse side effects induced by chemotherapeutic agents and radiation therapy have become of major importance to the clinical management of cancer patients.

20 Colorectal Cancer

Survival from colorectal cancer depends on the stage and grade of the tumor, for example precursor adenomas to metastatic adenocarcinoma. Generally, colorectal cancer can be treated by surgically removing
25 the tumor, but overall survival rates remain between 45 and 60 percent. Colonic excision morbidity rates are fairly low and is generally associated with the anastomosis and not the extent of the removal of the tumor and local tissue. In patients with a high risk of
30 reoccurrence, however, chemotherapy has been incorporated into the treatment regimen in order to improve survival rates.

-5-

Tumor metastasis prior to surgery is generally believed to be the cause of surgical intervention failure and up to one year of chemotherapy is required to kill the non-excised tumor cells. As severe toxicity is associated with the chemotherapeutic agents, only patients at high risk of recurrence are placed on chemotherapy following surgery.

Prostate Cancer

Prostate cancer is now the leading form of cancer among men and the second most frequent cause of death from cancer in men. It is estimated that more than 165,000 new cases of prostate cancer were diagnosed in 1993, and more than 35,000 men died from prostate cancer in that year. Additionally, the incidence of prostate cancer has increased by 50% since 1981, and mortality from this disease has continued to increase. Previously, most men died of other illnesses or diseases before dying from their prostate cancer. We now face increasing morbidity from prostate cancer as men live longer and the disease has the opportunity to progress.

Current therapies for prostate cancer focus upon reducing levels of dihydrotestosterone to decrease or prevent growth of prostate cancer. Radiation alone or in combination with surgery and/or chemotherapeutic agents is often used.

In addition to the use of digital rectal examination and transrectal ultrasonography, prostate-specific antigen (PSA) concentration is frequently used in the diagnosis of prostate cancer.

U.S. Pat. No. 4,472,382 discloses treatment of benign prostatic hyperplasia (BPH) with an antiandrogen and certain peptides which act as LH-RH agonists. U.S.

-6-

Pat. No. 4,596,797 discloses aromatase inhibitors as a method of prophylaxis and/or treatment of prostatic hyperplasia. U.S. Pat. No. 4,760,053 describes a treatment of certain cancers which combines an LHRH agonist with an antiandrogen and/or an antiestrogen and/or at least one inhibitor of sex steroid biosynthesis. U.S. Pat. No. 4,775,660 discloses a method of treating breast cancer with a combination therapy which may include surgical or chemical prevention of ovarian secretions and administering an antiandrogen and an antiestrogen. U.S. Pat. No. 4,659,695 discloses a method of treatment of prostate cancer in susceptible male animals including humans whose testicular hormonal secretions are blocked by surgical or chemical means, e.g. by use of an LHRH agonist, which comprises administering an antiandrogen, e.g. flutamide, in association with at least one inhibitor of sex steroid biosynthesis, e.g. aminoglutethimide and/or ketoconazole.

20 Prostate Specific Antigen

One well known prostate cancer marker is Prostate Specific Antigen (PSA). PSA is a protein produced by prostate cells and is frequently present at elevated levels in the blood of men who have prostate cancer. PSA has been shown to correlate with tumor burden, serve as an indicator of metastatic involvement, and provide a parameter for following the response to surgery, irradiation, and androgen replacement therapy in prostate cancer patients. It should be noted that Prostate Specific Antigen (PSA) is a completely different protein from Prostate Specific Membrane Antigen (PSMA). The two proteins have different

-7-

structures and functions and should not be confused because of their similar nomenclature.

Prostate Specific Membrane Antigen (PSMA)

In 1993, the molecular cloning of a prostate-specific membrane antigen (PSMA) was reported as a potential prostate carcinoma marker and hypothesized to serve as a target for imaging and cytotoxic treatment modalities for prostate cancer. Antibodies against PSMA have been described and examined clinically for diagnosis and treatment of prostate cancer. In particular, Indium-111 labeled PSMA antibodies have been described and examined for diagnosis of prostate cancer and indium-labeled PSMA antibodies have been described and examined for the treatment of prostate cancer.

Pancreas Cancer

Approximately 2% of new cancer cases diagnoses in the United States is pancreatic cancer. Pancreatic cancer is generally classified into two clinical types: 1) adenocarcinoma (metastatic and non-metastatic), and 2) cystic neoplasms (serous cystadenomas, mucinous cystic neoplasms, papillary cystic neoplasms, acinar cell cystadenocarcinoma, cystic choriocarcinoma, cystic teratomas, angiomatous neoplasms).

Ovary Cancer

Celomic epithelial carcinoma accounts for approximately 90% of ovarian cancer cases. Preferred single agents that can be used in combination include: alkylating agents, ifosfamide, cisplatin, carboplatin, taxol, doxorubicin, 5-fluorouracil, methotrexate, mitomycin, hexamethylmelamine, progestins, antiestrogens, prednimustine, dihydroxybusulfan, galactitol, interferon alpha and interferon gamma.

-8-

Cancer of the fallopian tube is the least common type of ovarian cancer, accounting for approximately 400 new cancer cases per year in the United States.

Papillary serous adenocarcinoma accounts for
5 approximately 90% of all malignancies of the ovarian tube.

Detailed Description of the Invention

Treatment of a neoplasia disorder in a mammal in
10 need of such treatment is provided by methods and combinations using radiation and a MMP inhibitor. The method comprises treating a mammal with a therapeutically effective amount of a combination comprising a MMP inhibitor and a radiotherapeutic agent.
15 Besides being useful for human treatment, the present invention is also useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.
20 Inhibitors of MMP potentiate tumor response to radiation. Thus, MMP inhibitors improve the efficacy of radiotherapy.

The methods and combinations of the present invention may be used for the treatment of neoplasia
25 disorders selected from the group consisting of acral lentiginous melanoma, actinic keratoses, adenocarcinoma, adenoid cystic carcinoma, adenomas, adenosarcoma, adenosquamous carcinoma, astrocytic tumors, bartholin gland carcinoma, basal cell carcinoma, bronchial gland
30 carcinomas, capillary, carcinoids, carcinoma, carcinosarcoma, cavernous, cholangiocarcinoma, chondrosarcoma, choroid plexus papilloma/carcinoma,

-9-

clear cell carcinoma, cystadenoma, endodermal sinus tumor, endometrial hyperplasia, endometrial stromal sarcoma, endometrioid adenocarcinoma, ependymal, epitheloid, Ewing's sarcoma, fibrolamellar, focal
5 nodular hyperplasia, gastrinoma, germ cell tumors, glioblastoma, glucagonoma, hemangiblastomas, hemangioendothelioma, hemangiomas, hepatic adenoma, hepatic adenomatosis, hepatocellular carcinoma, insulinoma, intraepithelial neoplasia, interepithelial
10 squamous cell neoplasia, invasive squamous cell carcinoma, large cell carcinoma, leiomyosarcoma, lentigo maligna melanomas, malignant melanoma, malignant mesothelial tumors, medulloblastoma, medulloepithelioma, melanoma, meningeal, mesothelial, metastatic carcinoma,
15 mucoepidermoid carcinoma, neuroblastoma, neuroepithelial adenocarcinoma nodular melanoma, oat cell carcinoma, oligodendroglial, osteosarcoma, pancreatic polypeptide, papillary serous adenocarcinoma, pineal cell, pituitary tumors, plasmacytoma, pseudosarcoma, pulmonary blastoma,
20 renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, serous carcinoma, small cell carcinoma, soft tissue carcinomas, somatostatin-secreting tumor, squamous carcinoma, squamous cell carcinoma, submesothelial, superficial spreading melanoma,
25 undifferentiated carcinoma, uveal melanoma, verrucous carcinoma, vipoma, well differentiated carcinoma, and Wilm's tumor.

The methods and compositions of the present invention provide one or more benefits. A combination
30 of a MMP inhibitor with radiation therapy of the present invention are useful in treating neoplasia disorders. Preferably, the MMP inhibitor agent or agents and the

-10-

radiation therapies of the present invention is administered in combination at a low dose, that is, at a dose lower than has been conventionally used in clinical situations for each of the individual components administered alone.

A benefit of lowering the dose of the radiation therapies of the present invention administered to a mammal includes a decrease in the incidence of adverse effects associated with higher dosages.

By lowering the incidence of adverse effects, an improvement in the quality of life of a patient undergoing treatment for cancer is contemplated. Further benefits of lowering the incidence of adverse effects include an improvement in patient compliance, and a reduction in the number of hospitalizations needed for the treatment of adverse effects.

Alternatively, the methods and combination of the present invention can also maximize the therapeutic effect at higher doses.

The term "pharmaceutically acceptable" is used herein to mean that the modified noun is appropriate for use in a pharmaceutical product. Pharmaceutically acceptable cations include metallic ions and organic ions. More preferred metallic ions include, but are not limited to appropriate alkali metal salts, alkaline earth metal salts and other physiological acceptable metal ions. Exemplary ions include aluminum, calcium, lithium, magnesium, potassium, sodium and zinc in their usual valences. Preferred organic ions include protonated tertiary amines and quaternary ammonium cations, including in part, trimethylamine, diethylamine, N,N'-dibenzylethylenediamine,

-11-

chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. Exemplary pharmaceutically acceptable acids include without limitation hydrochloric acid,

5 hydrobromic acid, phosphoric acid, sulfuric acid, methanesulfonic acid, acetic acid, formic acid, tartaric acid, maleic acid, malic acid, citric acid, isocitric acid, succinic acid, lactic acid, gluconic acid, glucuronic acid, pyruvic acid oxalacetic acid, fumaric

10 acid, propionic acid, aspartic acid, glutamic acid, benzoic acid, and the like.

Also included in the combination of the invention are the isomeric forms and tautomers of the described compounds and the pharmaceutically-acceptable salts

15 thereof. Illustrative pharmaceutically acceptable salts are prepared from formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic,

20 stearic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, cyclohexylaminosulfonic, algenic,

25 β -hydroxybutyric, galactaric and galacturonic acids.

Suitable pharmaceutically-acceptable base addition salts of compounds of the present invention include metallic ion salts and organic ion salts. More preferred metallic ion salts include, but are not

30 limited to appropriate alkali metal (group Ia) salts, alkaline earth metal (group IIa) salts and other

-12-

physiological acceptable metal ions. Such salts can be made from the ions of aluminum, calcium, lithium, magnesium, potassium, sodium and zinc. Preferred organic salts can be made from tertiary amines and quaternary ammonium salts, including in part, trimethylamine, diethylamine, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of the above salts can be prepared by those skilled in the art by conventional means from the corresponding compound of the present invention.

A MMP inhibitor of the present invention can be formulated as a pharmaceutical composition. Such a composition can then be administered orally, parenterally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration can also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques. Formulation of drugs is discussed in, for example, Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania; 1975 and Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable

-13-

dispersing or wetting agents and suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. Dimethyl acetamide, surfactants including ionic and non-ionic detergents, polyethylene glycols can be used. Mixtures of solvents and wetting agents such as those discussed above are also useful.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter, synthetic mono- di- or triglycerides, fatty acids and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration can include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, a contemplated aromatic sulfone hydroximate inhibitor compound can be admixed with lactose, sucrose, starch powder, cellulose

-14-

esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate,

5 polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets can contain a controlled-release formulation as can be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. In
10 the case of capsules, tablets, and pills, the dosage forms can also comprise buffering agents such as sodium citrate, magnesium or calcium carbonate or bicarbonate. Tablets and pills can additionally be prepared with enteric coatings.

15 For therapeutic purposes, formulations for parenteral administration can be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions can be prepared from sterile powders or granules having one or
20 more of the carriers or diluents mentioned for use in the formulations for oral administration. A contemplated MMP inhibitor compound can be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl
25 alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions,
30 solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions can also comprise adjuvants, such as

-15-

wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the mammalian host treated and the particular mode of administration.

The term "treatment" refers to any process, action, application, therapy, or the like, wherein a mammal, including a human being, is subject to medical aid with the object of improving the mammal's condition, directly or indirectly.

The term "inhibition," in the context of neoplasia, tumor growth or tumor cell growth, may be assessed by delayed appearance of primary or secondary tumors, slowed development of primary or secondary tumors, decreased occurrence of primary or secondary tumors, slowed or decreased severity of secondary effects of disease, arrested tumor growth and regression of tumors, among others. In the extreme, complete inhibition, is referred to herein as prevention.

The phrase "combination therapy" (or "co-therapy") embraces the administration of a matrix metalloproteinase inhibitor and radiation therapy, and, optionally, an antineoplastic agent, as part of a specific treatment regimen intended to provide a beneficial effect from the co-action of the matrix metalloproteinase inhibitor and the radiation therapy. The beneficial effect of the combination includes, but is not limited to, pharmacokinetic or pharmacodynamic co-action resulting from the combination of the matrix metalloproteinase inhibitor and the radiation therapy. Administration of the matrix metalloproteinase inhibitor

-16-

and the radiation therapy in combination typically is carried out over a defined time period (usually minutes, hours, days or weeks depending upon the combination selected). "Combination therapy" generally is not intended to encompass the administration of a matrix metalloproteinase inhibitor and radiation therapy as part of separate monotherapy regimens that incidentally and arbitrarily result in the combinations of the present invention. "Combination therapy" is intended to embrace administration of a matrix metalloproteinase inhibitor and radiation therapy in a sequential manner, that is, wherein the matrix metalloproteinase inhibitor and the radiation therapy are administered at different times, as well as administration of the matrix metalloproteinase and radiation therapy in a substantially simultaneous manner. Substantially simultaneous administration can be accomplished, for example, by administering to the subject concurrently with radiation therapy a single capsule having a fixed ratio of each therapeutic agent or in multiple, single capsules for each therapeutic agent. Sequential or substantially simultaneous administration of each therapeutic agent can be effected by any appropriate route including, but not limited to, oral routes, intravenous routes, intramuscular routes, and direct absorption through mucous membrane tissues. The therapeutic agents, if more than one, can be administered by the same route or by different routes. For example, a first therapeutic agent of the combination selected may be administered by intravenous injection while the other therapeutic agents of the combination may be administered orally. Alternatively,

-17-

for example, all therapeutic agents may be administered orally or all therapeutic agents may be administered by intravenous injection. The sequence in which the matrix metalloproteinase inhibitor and radiation therapy are

5 administered is not narrowly critical although radiation therapy typically will follow the administration of the matrix metalloproteinase inhibitor. "Combination therapy" also can embrace the administration of the matrix metalloproteinase inhibitor and radiation therapy

10 as described above in further combination with other biologically active ingredients (such as, but not limited to, an antineoplastic agent) and non-drug therapies (such as, but not limited to, surgery). The radiation treatment of the combination may be conducted

15 at any suitable time so long as a beneficial effect from the co-action of the combination of the matrix metalloproteinase inhibitor and radiation treatment is achieved. For example, in appropriate cases, the beneficial effect is still achieved even when the

20 radiation treatment is temporally removed from the administration of the matrix metalloproteinase inhibitor, perhaps by days or even weeks.

The term "prevention" includes either preventing the onset of clinically evident neoplasia altogether or

25 preventing the onset of a preclinically evident stage of neoplasia in individuals at risk. Also intended to be encompassed by this definition is the prevention of initiation for malignant cells or to arrest or reverse the progression of premalignant cells to malignant

30 cells. This includes prophylactic treatment of those at risk of developing the neoplasia.

-18-

Angiogenesis is an attractive therapeutic target because it is a multi-step process that occurs in a specific sequence, thus providing several possible targets for drug action. Examples of agents that
5 interfere with several of these steps include specific MMP inhibitors.

The phrase "therapeutically-effective" is intended to qualify the amount of each agent that will achieve the goal of improvement in neoplastic disease severity
10 and the frequency of incidence over treatment of each agent by itself, while avoiding adverse side effects typically associated with alternative therapies.

A "therapeutic effect" relieves to some extent one or more of the symptoms of a neoplasia disorder. In
15 reference to the treatment of a cancer, a therapeutic effect refers to one or more of the following: 1) reduction in the number of cancer cells; 2) reduction in tumor size; 3) inhibition (i.e., slowing to some extent, preferably stopping) of cancer cell infiltration into
20 peripheral organs; 4) inhibition (i.e., slowing to some extent, preferably stopping) of tumor metastasis; 5) inhibition, to some extent, of tumor growth; 6) relieving or reducing to some extent one or more of the symptoms associated with the disorder; and/or 7)
25 relieving or reducing the side effects associated with the administration of anticancer agents.

"Therapeutic effective amount" is intended to qualify the amount required to achieve a therapeutic effect.

30 The phrases "low dose" or "low dose amount", in characterizing a therapeutically effective amount of the MMP inhibitor and the radiation or therapy in the

combination therapy, defines a quantity of such therapy, or a range of quantity of such therapy, that is capable of diminishing the neoplastic disease while reducing or avoiding one or more radiation-induced side effects, such as myelosuppression, cardiac toxicity, skin erythema and desquamation, alopecia, inflammation or fibrosis.

The phrase "adjunctive therapy" includes agents such as those, for example, that reduce the toxic effect of anticancer drugs, e.g., bone resorption inhibitors, cardioprotective agents; prevent or reduce the incidence of nausea and vomiting associated with chemotherapy, radiotherapy or operation; or reduce the incidence of infection associated with the administration of myelosuppressive anticancer drugs.

The phrase a "radiotherapeutic agent" refers to the use of electromagnetic or particulate radiation in the treatment of neoplasia. Examples of radiotherapeutic agents are provided in, but not limited to, radiation therapy and is known in the art (Hellman, Principles of Radiation Therapy, Cancer, in Principles and Practice of Oncology, 248-75 (Devita et al., ed., 4th edit., volume 1, 1993).

The term "clinical tumor" includes neoplasms that are identifiable through clinical screening or diagnostic procedures including, but not limited to, palpation, biopsy, cell proliferation index, endoscopy, mammography, digital mammography, ultrasonography, computed tomography (CT), magnetic resonance imaging (MRI), positron emission tomography (PET), radiography, radionuclide evaluation, CT- or MRI-guided aspiration cytology, and imaging-guided needle biopsy, among others. Such diagnostic techniques are well known to

those skilled in the art and are described in Cancer
Medicine 4th Edition, Volume One. J.F. Holland, R.C.
Bast, D.L. Morton, E. Frei III, D.W. Kufe, and R.R.
Weichselbaum (Editors). Williams & Wilkins, Baltimore
5 (1997).

The term "tumor marker" or "tumor biomarker"
encompasses a wide variety of molecules with divergent
characteristics that appear in body fluids or tissue in
association with a clinical tumor and also includes
10 tumor-associated chromosomal changes. Tumor markers fall
primarily into three categories: molecular or cellular
markers, chromosomal markers, and serological or serum
markers. Molecular and chromosomal markers complement
standard parameters used to describe a tumor (i.e.
15 histopathology, grade, tumor size) and are used
primarily in refining disease diagnosis and prognosis
after clinical manifestation. Serum markers can often
be measured many months before clinical tumor detection
and are thus useful as an early diagnostic test, in
20 patient monitoring, and in therapy evaluation.

Molecular Tumor Markers

Molecular markers of cancer are products of cancer
cells or molecular changes that take place in cells
because of activation of cell division or inhibition of
25 apoptosis. Expression of these markers can predict a
cell's malignant potential. Because cellular markers
are not secreted, tumor tissue samples are generally
required for their detection. Non-limiting examples of
molecular tumor markers that can be used in the present
30 invention are listed in Table No. 1, below.

Table No. 1. Non-limiting Examples of Molecular Tumor
Markers

Tumor	Marker
Breast	p53
Breast, Ovarian	ErbB-2/Her-2
Breast	S phase and ploidy
Breast	pS2
Breast	MDR2
Breast	urokinase plasminogen activator
Breast, Colon, Lung	myc family

Chromosomal Tumor Markers

Somatic mutations and chromosomal aberrations have been associated with a variety of tumors. Since the

5 identification of the Philadelphia Chromosome by Nowel and Hungerford, a wide effort to identify tumor-specific chromosomal alterations has ensued. Chromosomal cancer markers, like cellular markers, are can be used in the diagnosis and prognosis of cancer. In addition to the

10 diagnostic and prognostic implications of chromosomal alterations, it is hypothesized that germ-line mutations can be used to predict the likelihood that a particular person will develop a given type of tumor. Non-limiting examples of chromosomal tumor markers that can be used

15 in the present invention are listed in Table No. 2, below.

Table No. 2. Non-limiting Examples of Chromosomal Tumor Markers

Tumor	Marker
Breast	1p36 loss

-22-

Breast	6q24-27 loss
Breast	11q22-23 loss
Breast	11q13 amplification
Breast	TP53 mutation
Colon	Gain of chromosome 13
Colon	Deletion of short arm of chromosome 1
Lung	Loss of 3p
Lung	Loss of 13q
Lung	Loss of 17p
Lung	Loss of 9p

Serological Tumor Markers

Serum markers including soluble antigens, enzymes and hormones comprise a third category of tumor markers.

5 Monitoring serum tumor marker concentrations during therapy provides an early indication of tumor recurrence and of therapy efficacy. Serum markers are advantageous for patient surveillance compared to chromosomal and cellular markers because serum samples are more easily

10 obtainable than tissue samples, and because serum assays can be performed serially and more rapidly. Serum tumor markers can be used to determine appropriate therapeutic doses within individual patients. For example, the efficacy of a combination regimen consisting of

15 chemotherapeutic and antiangiogenic agents can be measured by monitoring the relevant serum cancer marker levels. Moreover, an efficacious therapy dose can be achieved by modulating the therapeutic dose so as to keep the particular serum tumor marker concentration

20 stable or within the reference range, which may vary depending upon the indication. The amount of therapy

can then be modulated specifically for each patient so as to minimize side effects while still maintaining stable, reference range tumor marker levels. Table No. 3 provides non-limiting examples of serological tumor markers that can be used in the present invention.

Table No. 3. Non-limiting Examples of Serum Tumor Markers

Cancer Type	Marker
Germ Cell Tumors	a-fetoprotein (AFP)
Germ Cell Tumors	human chorionic gonadotrophin (hCG)
Germ Cell Tumors	placental alkaline phosphatase (PLAP)
Germ Cell Tumors	lactate dehydrogenase (LDH)
Prostate	prostate specific antigen (PSA)
Breast	carcinoembryonic antigen (CEA)
Breast	MUC-1 antigen (CA15-3)
Breast	tissue polypeptide antigen (TPA)
Breast	tissue polypeptide specific antigen (TPS)
Breast	CYFRA 21.1
Breast	soluble erb-B-2
Ovarian	CA125
Ovarian	OVX1
Ovarian	cancer antigen CA72-4
Ovarian	TPA

-24-

Ovarian	TPS
Gastrointestinal	CD44v6
Gastrointestinal	CEA
Gastrointestinal	cancer antigen CA19-9
Gastrointestinal	NCC-ST-439 antigen (Dukes C)
Gastrointestinal	cancer antigen CA242
Gastrointestinal	soluble <i>erb</i> -B-2
Gastrointestinal	cancer antigen CA195
Gastrointestinal	TPA
Gastrointestinal	YKL-40
Gastrointestinal	TPS
Esophageal	CYFRA 21-1
Esophageal	TPA
Esophageal	TPS
Esophageal	cancer antigen CA19-9
Gastric Cancer	CEA
Gastric Cancer	cancer antigen CA19-9
Gastric Cancer	cancer antigen CA72-4
Lung	neruon specific enolase (NSE)
Lung	CEA
\Lung	CYFRA 21-1
Lung	cancer antigen CA 125
Lung	TPA
Lung	squamous cell carcinoma antigen (SCC)
Pancreatic cancer	ca19-9
Pancreatic cancer	ca50
Pancreatic cancer	ca119
Pancreatic cancer	ca125
Pancreatic cancer	CEA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

-25-

Pancreatic cancer	
Renal Cancer	CD44v6
Renal Cancer	E-cadherin
Renal Cancer	PCNA (proliferating cell nuclear antigen)

Examples

Germ Cell Cancers

Non-limiting examples of tumor markers useful in the present invention for the detection of germ cell cancers include, but are not limited to, a-fetoprotein (AFP), human chorionic gonadotrophin (hCG) and its beta subunit (hCGb), lactate dehydrogenase (LDH), and placental alkaline phosphatase (PLAP).

AFP has an upper reference limit of approximately -10 kU/L after the first year of life and may be elevated in germ cell tumors, hepatocellular carcinoma and also in gastric, colon, biliary, pancreatic and lung cancers. AFP serum half life is approximately five days after orchidectomy. According to EGTM recommendations, AFP serum levels less than 1,000 kU/L correlate with a good prognosis, AFP levels between 1,000 and 10,000 kU/L, inclusive, correlate with intermediate prognosis, and AFP levels greater than 10,000 U/L correlate with a poor prognosis.

HCG is synthesized in the placenta and is also produced by malignant cells. Serum hCG concentrations may be increased in pancreatic adenocarcinomas, islet cell tumors, tumors of the small and large bowel, hepatoma, stomach, lung, ovaries, breast and kidney. Because some tumors only hCGb, measurement of both hCG

-26-

and hCGb is recommended. Normally, serum hCG in men and pre-menopausal women is as high as -5 U/L while post-menopausal women have levels up to -10 U/L. Serum half life of hCG ranges from 16-24 hours. According to the

5 EGTM, hCG serum levels under 5000 U/L correlate with a good prognosis, levels between 5000 and 50000 U/L, inclusively correlate with an intermediate prognosis, and hCG serum levels greater than 50000 U/L correlate with a poor prognosis. Further, normal hCG half lives

10 correlate with good prognosis while prolonged half lives correlate with poor prognosis.

LDH is an enzyme expressed in cardiac and skeletal muscle as well as in other organs. The LDH-1 isoenzyme is most commonly found in testicular germ cell tumors

15 but can also occur in a variety of benign conditions such as skeletal muscle disease and myocardial infarction. Total LDH is used to measure independent prognostic value in patients with advanced germ cell tumors. LDH levels less than 1.5 x the reference range

20 are associated with a good prognosis, levels between 1.5 and 10 x the reference range, inclusive, are associated with an intermediate prognosis, and levels more than 10 x the reference range are associated with a poor prognosis.

25 PLAP is a enzyme of alkaline phosphatase normally expressed by placental syncytiotrophoblasts. Elevated serum concentrations of PLAP are found in seminomas, non-seminomatous tumors, and ovarian tumors, and may also provide a marker for testicular tumors. PLAP has a

30 normal half life after surgical resection of between 0.6 and 2.8 days.

Prostate Cancer

-27-

A nonlimiting example of a tumor marker useful in the present invention for the detection of prostate cancer is prostate specific antigen (PSA). PSA is a glycoprotein that is almost exclusively produced in the prostate. In human serum, uncomplexed f-PSA and a complex of f-PSA with α 1-antichymotrypsin make up total PSA (t-PSA). T-PSA is useful in determining prognosis in patients that are not currently undergoing anti-androgen treatment. Rising t-PSA levels via serial measurement indicate the presence of residual disease.

Breast Cancer

Non-limiting examples of serum tumor markers useful in the present invention for the detection of breast cancer include, but is not limited to carcinoembryonic antigen (CEA) and MUC-1 (CA 15.3). Serum CEA and CA15.3 levels are elevated in patients with node involvement compared to patients without node involvement, and in patients with larger tumors compared to smaller tumors. Normal range cutoff points (upper limit) are 5-10 mg/L for CEA and 35-60 u/ml for CA15.3. Additional specificity (99.3%) is gained by confirming serum levels with two serial increases of more than 15%.

Ovarian Cancer

A non-limiting example of a tumor marker useful in the present invention for the detection of ovarian cancer is CA125. Normally, women have serum CA125 levels between 0-35 kU/L; 99% of post-menopausal women have levels below 20 kU/L. Serum concentration of CA125 after chemotherapy is a strong predictor of outcome as elevated CA125 levels are found in roughly 80% of all patients with epithelial ovarian cancer. Further, prolonged CA125 half-life or a less than 7-fold decrease

-28-

during early treatment is also a predictor of poor disease prognosis.

Gastrointestinal Cancers

A non-limiting example of a tumor marker useful in the present invention for the detection of colon cancer is carcinoembryonic antigen (CEA). CEA is a glycoprotein produced during embryonal and fetal development and has a high sensitivity for advanced carcinomas including those of the colon, breast, stomach and lung. High pre- or postoperative concentrations (>2.5 ng/ml) of CEA are associated with worse prognosis than are low concentrations. Further, some studies in the literature report that slow rising CEA levels indicates local recurrence while rapidly increasing levels suggests hepatic metastasis.

Lung Cancer

Examples of serum markers useful in the present invention to monitor lung cancer therapy include, but are not limited to, CEA, cytokeratin 19 fragments (CYFRA 21-1), and Neuron Specific Enolase (NSE).

NSE is a glycolytic isoenzyme of enolase produced in central and peripheral neurons and malignant tumors of neuroectodermal origin. At diagnosis, NSE concentrations greater than 25 ng/mL are suggestive of malignancy and lung cancer while concentrations greater than 100 ng/mL are suggestive of small cell lung cancer.

CYFRA 21-1 is a tumor marker test which uses two specific monoclonal antibodies against a cytokeratin 19 fragment. At diagnosis, CYFRA 21-1 concentrations greater than 10 ng/mL are suggestive of malignancy while concentrations greater than 30 ng/mL are suggestive of lung cancer.

-29-

Accordingly, dosing of the matrix metalloproteinase inhibitor and radiation therapy may be determined and adjusted based on measurement of tumor markers in body fluids or tissues, particularly based on tumor markers in serum. For example, a decrease in serum marker level relative to baseline serum marker prior to administration of the matrix metalloproteinase inhibitor and radiation therapy indicates a decrease in cancer-associated changes and provides a correlation with inhibition of the cancer. In one embodiment, therefore, the method of the present invention comprises administering the matrix metalloproteinase inhibitor and radiation therapy at doses that in combination result in a decrease in one or more tumor markers, particularly a decrease in one or more serum tumor markers, in the mammal relative to baseline tumor marker levels.

Similarly, decreasing tumor marker concentrations or serum half lives after administration of the combination indicates a good prognosis, while tumor marker concentrations which decline slowly and do not reach the normal reference range predict residual tumor and poor prognosis. Further, during follow-up therapy, increases in tumor marker concentration predicts recurrent disease many months before clinical manifestation.

In addition to the above examples, Table No. 4, below, lists several references, hereby individually incorporated by reference herein, that describe tumor markers and their use in detecting and monitoring tumor growth and progression.

The phrase "matrix metalloproteinase inhibitor" or "MMP inhibitor" includes agents that specifically inhibit a class of enzymes, the zinc metalloproteinases (metalloproteases). The zinc metalloproteinases are involved in the degradation of connective tissue or connective tissue components. These enzymes are released from resident tissue cells and/or invading inflammatory or tumor cells. Blocking the action of zinc metalloproteinases interferes with the creation of paths for newly forming blood vessels to follow. Examples of MMP inhibitors are described in Golub, LM, Inhibition of Matrix Metalloproteinases: Therapeutic Applications (Annals of the New York Academy of Science, Vol 878). Robert A. Greenwald and Stanley Zucker (Eds.), June 1999), and is hereby incorporated by reference.

Connective tissue, extracellular matrix constituents and basement membranes are required components of all mammals. These components are the biological materials that provide rigidity, differentiation, attachments and, in some cases, elasticity to biological systems including human beings and other mammals. Connective tissues components include, for example, collagen, elastin, proteoglycans, fibronectin and laminin. These biochemicals makeup, or are components of structures, such as skin, bone, teeth, tendon, cartilage, basement membrane, blood vessels, cornea and vitreous humor.

Under normal conditions, connective tissue turnover and/or repair processes are controlled and in equilibrium. The loss of this balance for whatever reason leads to a number of disease states. Inhibition of the enzymes responsible loss of equilibrium provides

a control mechanism for this tissue decomposition and, therefore, a treatment for these diseases.

Degradation of connective tissue or connective tissue components is carried out by the action of
5 proteinase enzymes released from resident tissue cells and/or invading inflammatory or tumor cells. A major class of enzymes involved in this function are the zinc metalloproteinases (metalloproteases).

The metalloprotease enzymes are divided into
10 classes with some members having several different names in common use. Examples are: collagenase I (MMP-1, fibroblast collagenase; EC 3.4.24.3); collagenase II (MMP-8, neutrophil collagenase; EC 3.4.24.34), collagenase III (MMP-13), stromelysin 1 (MMP-3; EC
15 3.4.24.17), stromelysin 2 (MMP-10; EC 3.4.24.22), proteoglycanase, matrilysin (MMP-7), gelatinase A (MMP-2, 72kDa gelatinase, basement membrane collagenase; EC 3.4.24.24), gelatinase B (MMP-9, 92kDa gelatinase; EC 3.4.24.35), stromelysin 3 (MMP-11), metalloelastase
20 (MMP-12, HME, human macrophage elastase) and membrane MMP (MMP-14). MMP is an abbreviation or acronym representing the term Matrix Metalloprotease with the attached numerals providing differentiation between specific members of the MMP group.

25 The uncontrolled breakdown of connective tissue by metalloproteases is a feature of many pathological conditions. Examples include rheumatoid arthritis, osteoarthritis, septic arthritis; corneal, epidermal or gastric ulceration; tumor metastasis, invasion or
30 angiogenesis; periodontal disease; proteinuria; Alzheimer's Disease; coronary thrombosis and bone disease. Defective injury repair processes also occur.

This can produce improper wound healing leading to weak repairs, adhesions and scarring. These latter defects can lead to disfigurement and/or permanent disabilities as with post-surgical adhesions.

- 5 Matrix metalloproteases are also involved in the biosynthesis of tumor necrosis factor (TNF) and inhibition of the production or action of TNF and related compounds is an important clinical disease treatment mechanism. TNF- α , for example, is a cytokine
- 10 that at present is thought to be produced initially as a 28 kD cell-associated molecule. It is released as an active, 17 kD form that can mediate a large integer of deleterious effects *in vitro* and *in vivo*. For example, TNF can cause and/or contribute to the effects of
- 15 inflammation, rheumatoid arthritis, autoimmune disease, multiple sclerosis, graft rejection, fibrotic disease, cancer, infectious diseases, malaria, mycobacterial infection, meningitis, fever, psoriasis, cardiovascular/pulmonary effects such as post-ischemic
- 20 reperfusion injury, congestive heart failure, hemorrhage, coagulation, hyperoxic alveolar injury, radiation damage and acute phase responses like those seen with infections and sepsis and during shock such as septic shock and hemodynamic shock. Chronic release of
- 25 active TNF can cause cachexia and anorexia. TNF can be lethal.

TNF- α convertase is a metalloproteinase involved in the formation of active TNF- α . Inhibition of TNF- α convertase inhibits production of active TNF- α .

- 30 Compounds that inhibit both MMPs activity have been disclosed in, for example PCT Publication WO 94/24140. Other compounds that inhibit both MMPs activity have

also been disclosed in WO 94/02466. Still other compounds that inhibit both MMPs activity have been disclosed in WO 97/20824.

There remains a need for effective MMP and TNF- α convertase inhibiting agents. Compounds that inhibit MMPs such as collagenase, stromelysin and gelatinase have been shown to inhibit the release of TNF (Gearing et al. *Nature* 376, 555-557 (1994)). McGeehan et al., *Nature* 376, 558-561 (1994) also reports such findings.

MMPs are involved in other biochemical processes in mammals as well. Included is the control of ovulation, post-partum uterine involution, possibly implantation, cleavage of APP (β -Amyloid Precursor Protein) to the amyloid plaque and inactivation of α_1 -protease inhibitor (α_1 -PI). Inhibition of these metalloproteases permits the control of fertility and the treatment or prevention of Alzheimers Disease. In addition, increasing and maintaining the levels of an endogenous or administered serine protease inhibitor drug or biochemical such as α_1 -PI supports the treatment and prevention of diseases such as emphysema, pulmonary diseases, inflammatory diseases and diseases of aging such as loss of skin or organ stretch and resiliency.

Inhibition of selected MMPs can also be desirable in other instances. Treatment of cancer and/or inhibition of metastasis and/or inhibition of angiogenesis are examples of approaches to the treatment of diseases wherein the selective inhibition of stromelysin (MMP-3), gelatinase (MMP-2), or collagenase III (MMP-13) are the relatively most important enzyme or enzymes to inhibit especially when compared with

collagenase I (MMP-1). A drug that does not inhibit collagenase I can have a superior therapeutic profile.

Inhibitors of metalloproteases are known. Examples include natural biochemicals such as tissue inhibitor of metalloproteinase (TIMP), α_2 -macroglobulin and their analogs or derivatives. These are high molecular weight protein molecules that form inactive complexes with metalloproteases. An integer of smaller peptide-like compounds that inhibit metalloproteases have been described. Mercaptoamide peptidyl derivatives have shown ACE inhibition *in vitro* and *in vivo*. Angiotensin converting enzyme (ACE) aids in the production of angiotensin II, a potent pressor substance in mammals and inhibition of this enzyme leads to the lowering of blood pressure.

Thiol group-containing amide or peptidyl amide-based metalloprotease (MMP) inhibitors are known as is shown in, for example, WO 95/12389. Thiol group-containing amide or peptidyl amide-based metalloprotease (MMP) inhibitors are also shown in WO 96/11209. Still further Thiol group-containing amide or peptidyl amide-based metalloprotease (MMP) inhibitors are shown in U.S. Patent No. 4,595,700. Hydroxamate group-containing MMP inhibitors are disclosed in a number of published patent applications that disclose carbon back-boned compounds, such as in WO 95/29892. Other published patents include WO 97/24117. Additionally, EP 0 780 386 further discloses hydroxamate group-containing MMP inhibitors. WO 90/05719 disclose hydroxamates that have a peptidyl back-bones or peptidomimetic back-bones. WO 93/20047 also discloses hydroxamates that have a peptidyl back-bones or peptidomimetic back-bones. Additionally, WO

95/09841 discloses disclose hydroxamates that have peptidyl back-bones or peptidomimetic back-bones. And WO 96/06074 further discloses hydroxamates that have peptidyl back-bones or peptidomimetic back-bones.

- 5 Schwartz et al., *Progr. Med. Chem.*, 29:271-334(1992) also discloses disclose hydroxamates that have peptidyl back-bones or peptidomimetic back-bones. Furthermore, Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997) discloses hydroxamates that have peptidyl back-bones or
10 peptidomimetic back-bones. Also, Denis et al., *Invest. New Drugs*, 15(3): 175-185 (1997) discloses hydroxamates that have a peptidyl back-bones or peptidomimetic back-bones as well.

- One possible problem associated with known MMP
15 inhibitors is that such compounds often exhibit the same or similar inhibitory effects against each of the MMP enzymes. For example, the peptidomimetic hydroxamate known as batimastat is reported to exhibit IC₅₀ values of about 1 to about 20 nanomolar (nM) against each of
20 MMP-1, MMP-2, MMP-3, MMP-7, and MMP-9. Marimastat, another peptidomimetic hydroxamate was reported to be another broad-spectrum MMP inhibitor with an enzyme inhibitory spectrum very similar to batimastat, except that marimastat exhibited an IC₅₀ value against MMP-3 of
25 230 nM. Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997).

- Meta analysis of data from Phase I/II studies using marimastat in patients with advanced, rapidly
progressive, treatment-refractory solid tumor cancers
30 (colorectal, pancreatic, ovarian, prostate), indicated a dose-related reduction in the rise of cancer-specific antigens used as surrogate markers for biological

activity. The most common drug-related toxicity of marimastat in those clinical trials was musculoskeletal pain and stiffness, often commencing in the small joints in the hands, spreading to the arms and shoulder. A short dosing holiday of 1-3 weeks followed by dosage reduction permits treatment to continue. Rasmussen et al., *Pharmacol. Ther.*, 75(1): 69-75 (1997). It is thought that the lack of specificity of inhibitory effect among the MMPs may be the cause of that effect.

In view of the importance of hydroxamate MMP inhibitor compounds in the treatment of several diseases and the lack of enzyme specificity exhibited by two of the more potent drugs now in clinical trials, it would be beneficial to use hydroxamates of greater enzyme specificity. This would be particularly the case if the hydroxamate inhibitors exhibited limited inhibition of MMP-1 that is relatively ubiquitous and as yet not associated with any pathological condition, while exhibiting quite high inhibitory activity against one or more of MMP-2, MMP-9 or MMP-13 that are associated with several pathological conditions.

Non-limiting examples of matrix metalloproteinase inhibitors that may be used in the present invention are identified in Table No. 5, below.

Table No. 5. Matrix metalloproteinase inhibitors.

Compound	Trade Name	Reference	Dosage
Biphenyl hydroxamate		WO 97/18188	
	AG-3067 (Agouron)	Winter Conf. Med. Bio-	

Compound	Trade Name	Reference	Dosage
	Pharm. Inc.)	organic Chem. 1997 January, 26- 31	
	AG-3340 (Agouron Pharm. Inc.)	WO 97/20824	50 mg/kg treatment of Lewis lung carcinomas in test animals
	AG-2024 (Agouron Pharm. Inc.)		
	AG-3365 (Agouron Pharm. Inc.)		
3(S)-N-hydroxy- 4-(4-[4- (imidazol-1- yl)phenoxy]benze nesulfonyl)-2,2- dimethyl- tetrahydro-2H- 1,4-thiazine-3- carboxamide, and derivatives thereof		WO 97/20824. FEBS (1992) 296 (3):263	In female Lewis rats, arthritis model: dose of 25 mg/kg/day gave 97.5% weight loss inhibition
Heteroaryl		WO 98/17643	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Compound	Trade Name	Reference	Dosage
succinamides derivatives			
	AG-3296 (Agouron Pharm. Inc.)		
	AG- 3287 (Agour on Pharm. Inc.)		
	AG-3293 (Agouron Pharm. Inc.)		
	AG-3294 (Agouron Pharm. Inc.)		
	AG-3067 (Agouron Pharm. Inc.)	Winter Conf Med Bio- organic Chem 1997 January 26-31	
2R,4S)-4- hydroxy-2- isobutyl-5- mercapto-N- [(1S)-2,2- dimethyl-1- methylcarbamoylp rotyl]		EP 0818443	

11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782
 783
 784
 785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796
 797
 798
 799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000
 1001
 1002
 1003
 1004
 1005
 1006
 1007
 1008
 1009
 1010
 1011
 1012
 1013
 1014
 1015
 1016
 1017
 1018
 1019
 1020
 1021
 1022
 1023
 1024
 1025
 1026
 1027
 1028
 1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048
 1049
 1050
 1051
 1052
 1053
 1054
 1055
 1056
 1057
 1058
 1059
 1060
 1061
 1062
 1063
 1064
 1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073
 1074
 1075
 1076
 1077
 1078
 1079
 1080
 1081
 1082
 1083
 1084
 1085
 1086
 1087
 1088
 1089
 1090
 1091
 1092
 1093
 1094
 1095
 1096
 1097
 1098
 1099
 1100
 1101
 1102
 1103
 1104
 1105
 1106
 1107
 1108
 1109
 1110
 1111
 1112
 1113
 1114
 1115
 1116
 1117
 1118
 1119
 1120
 1121
 1122
 1123
 1124
 1125
 1126
 1127
 1128
 1129
 1130
 1131
 1132
 1133
 1134
 1135
 1136
 1137
 1138
 1139
 1140
 1141
 1142
 1143
 1144
 1145
 1146
 1147
 1148
 1149
 1150
 1151
 1152
 1153
 1154
 1155
 1156
 1157
 1158
 1159
 1160
 1161
 1162
 1163
 1164
 1165
 1166
 1167
 1168
 1169
 1170
 1171
 1172
 1173
 1174
 1175
 1176
 1177
 1178
 1179
 1180
 1181
 1182
 1183
 1184
 1185
 1186
 1187
 1188
 1189
 1190
 1191
 1192
 1193
 1194
 1195
 1196
 1197
 1198
 1199
 1200
 1201
 1202
 1203
 1204
 1205
 1206
 1207
 1208
 1209
 1210
 1211
 1212
 1213
 1214
 1215
 1216
 1217
 1218
 1219
 1220
 1221
 1222
 1223
 1224
 1225
 1226
 1227
 1228
 1229
 1230
 1231
 1232
 1233
 1234
 1235
 1236
 1237
 1238
 1239
 1240
 1241
 1242
 1243
 1244
 1245
 1246
 1247
 1248
 1249
 1250
 1251
 1252
 1253
 1254
 1255
 1256
 1257
 1258
 1259
 1260
 1261
 1262
 1263
 1264
 1265
 1266
 1267
 1268
 1269
 1270
 1271
 1272
 1273
 1274
 1275
 1276
 1277
 1278
 1279
 1280
 1281
 1282
 1283
 1284
 1285
 1286
 1287
 1288
 1289
 1290
 1291
 1292
 1293
 1294
 1295
 1296
 1297
 1298
 1299
 1300
 1301
 1302
 1303
 1304
 1305
 1306
 1307
 1308
 1309
 1310
 1311
 1312
 1313
 1314
 1315
 1316
 1317
 1318
 1319
 1320
 1321
 1322
 1323
 1324
 1325
 1326
 1327
 1328
 1329
 1330
 1331
 1332
 1333
 1334
 1335
 1336
 1337
 1338
 1339
 1340
 1341
 1342
 1343
 1344
 1345
 1346
 1347
 1348
 1349
 1350
 1351
 1352
 1353
 1354
 1355
 1356
 1357
 1358
 1359
 1360
 1361
 1362
 1363
 1364
 1365
 1366
 1367
 1368
 1369
 1370
 1371
 1372
 1373
 1374
 1375
 1376
 1377
 1378
 1379
 1380
 1381
 1382
 1383
 1384
 1385
 1386
 1387
 1388
 1389
 1390
 1391
 1392
 1393
 1394
 1395
 1396
 1397
 1398
 1399
 1400
 1401
 1402
 1403
 1404
 1405
 1406
 1407
 1408
 1409
 1410
 1411
 1412
 1413
 1414
 1415
 1416
 1417
 1418
 1419
 1420
 1421
 1422
 1423
 1424
 1425
 1426
 1427
 1428
 1429
 1430
 1431
 1432
 1433
 1434
 1435
 1436
 1437
 1438
 1439
 14

Compound	Trade Name	Reference	Dosage
pentanamide			
N-alkyl, N-phenylsulfonyl-N'-hydroxamic acid derivatives of heteroaryl carboxylic acids		WO 98/16520	
Novel N-alkyl, N-phenylsulfonyl-N'-hydroxamic acid derivatives of heteroaryl carboxylic acids		WO 98/16514	
Novel N-alkyl, N-phenylsulfonyl-N'-hydroxamic acid derivatives of cycloalkane carboxylic acids		WO 98/16506	
Novel N-alkyl, N-phenylsulfonyl-N'-hydroxamic acid derivatives of anthranilic acid		WO 98/16503	
sulfonamido-hydroxamic acid derivatives		EP 03/98753	

Compound	Trade Name	Reference	Dosage
TIMP-3: polynucleotides encoding endogenous (human) peptides		WO 95/09918	
(3alpha, 5beta,6alpha,7a1 phabeta)-4',4'- (hexahydro-2,2- dimethyl-1,3- benzodioxole-5, 6-diyl)bis(2,6- piperazinedione) and derivatives thereof		WO 93/23075	
	BE-16627B	WO 91/08222. Int. J. Cancer 1994 58 5 730 - 735	
(2S)-4-(4-(4- chlorophenyl)phe nyl)-4-oxo- 2- (2- phthalimidoethyl)butanoic acid		WO 96/15096	
	Bay-12- 9566	WO 96/15096	10 to 400 mg/day
4-oxo-2-(2- phthalimidoethyl) alcanoic acid		WO 97/43238	

[illegible]

Compound	Trade Name	Reference	Dosage
analogs of marimastat		Med Chem Lett 1998 8 11 1359 - 1364	
	GI-245402 (BB-2983)		
Hydroxamic acid derivatives		WO 94/21625	
Succinyl hydroxamic acid, N-formyl-N- hydroxy amino carboxylic acid and succinic acid amide derivatives		WO 95/32944	
hydroxamic acid, N-formyl-N- hydroxyamino and carboxylic acid derivatives,		WO 97/19053	
pseudopeptide hydroxamic and carboxylic acid derivatives from the corresponding lactone and alpha-amino acid		WO 97/19050	
Succinic acid amide		WO 97/03966. GB 95/00111.	

Compound	Trade Name	Reference	Dosage
derivatives		GB 95/00121.	
Hydroxamic acid derivatives		WO 97/02239	
Succinamidyl (alpha substituted) hydroxamic acid derivatives		WO 96/33165	
(2S,3R)-3-[2,2-dimethyl-1S-(thiazol-2-ylcarbamoyl)propylcarbamoyl]-5-methyl-2-(prop-2-enyl)hexano-hydroxamic acid and derivatives thereof		WO 96/25156	
Hydroxamic or carboxylic acid derivatives		WO 96/16931	
hydroxamic and carboxylic acids		WO 96/06074	
2-[(1S)-1-((1R)-2-[[1,1'-biphenyl]-4-ylmethylthio]-1-[(1S)-2,2-dimethyl-1-(methylcarbamoyl)propylcarbamoyl		WO 98/23588	

Publ. No. WO 00/38717

Compound	Trade Name	Reference	Dosage
]ethylcarbamoyl)-4-(1,3-dioxo-1,3-dihydroisoindol-2-yl)butylthio]-acetate, and derivatives thereof			
Hydroxamic acid derivatives as inhibitors of cytokine production		WO 95/09841	
Hydroxamic acid derivatives		WO 94/24140	
Aromatic or heteroaryl substituted hydroxamic or carboxylic acid derivatives		WO 95/19956	
Hydroxamic acid derivatives		WO 95/19957	Doses are preferably 1 to 100 mg/kg.
Hydroxamic acid and carboxylic acid derivatives		WO 95/19961	Doses are preferably 1 to 100 mg/kg.
Butanediamide, N1-	BB-1433		At 50 mg/kg bid. p.o.

Compound	Trade Name	Reference	Dosage
[1(cyclohexyl-methyl)-2-(methylamino)-2-oxoethyl]-N4,3-dihydroxy-2-(2-methylpropyl)-, [2R[N1(S*),2R*,3S*]]-			inhibited bone mineral density loss
tetracycline analogs and D-penicillamine		EP 733369	D-penicillamine reduced allergic encephalitis symptom scores in a dose dependent manner at 27, 125 and 375 mug with complete inhibition
	CDP-845	Biochem Pharmacol 1990 39 12 2041-2049	
succinamide derivatives		WO 95/04033	oral bioavail-ability by murine

T0300T = 320000

Compound	Trade Name	Reference	Dosage
			pleural cavity assay in the presence of gelatinase: Between 73% and 100% inhibition was displayed at 10 mg/kg for six of the compounds. The seventh displayed 100% inhibition at 80 mg/kg.
Peptidyl derivatives		WO 94/25435. WO 94/25434	
Mercaptoalkyl-peptidyl compounds having an imidazole substituent		WO 97/19075	
mercaptoalkyl-peptide derivatives		WO 97/38007. WO 95/12389. WO 96/11209.	

Compound	Trade Name	Reference	Dosage
Mercaptoalkyl- amide derivatives		WO 97/37974	
arylsulfonyl- hydrazine derivatives		WO 97/37973. WO 95/12389	
N-acetylthio- lacetyl-N-(3- phthalimidopropy l)-L-leucyl-L- phenylalanine N- methyleamide		WO 96/35714	
2-acetylsulfany- 1-5-phthalimido- pentanoyl-L- leucineN-(2- phenylethyl)- amide		WO 96/35712	dosages of about 0.5 mg to 3.5 g per day for the treatment of inflam- mation
5-phthalimido- pentanoyl-L- leucyl-L- phenylalanineN- methyleamide		WO 96/35711	
peptidyl derivatives		WO 98/06696	
4-[4- (methoxycarbonyl methoxy)-3,5- dimethylphenyl]-		WO 98/05635	

Compound	Trade Name	Reference	Dosage
2-methyl-1(2H)-phthalazinone, and hydroxamic and carboxylic acid derivatives			
thio-substituted peptides		WO 97/12902	
Mercaptoamides		WO 97/12861	
Peptidyl derivatives having SH or acylo groups which are amides, primary amides or thioamides		WO 96/35687	
	D-5410 (Chiro-science Group plc)		
		WO 95/13289	
	CH-104, (Chiro-science Group plc)		
	D-2163 (Chiro Science Ltd.)		
	D-1927 (Chiro		

Publ. No. 99/0038717

Compound	Trade Name	Reference	Dosage
	Science Ltd.)		
	Dermastat (Colla- Genex Phar- maceu- tical Inc.)		
	Metastat (Colla- Genex)		
	Osteostat (Colla- Genex Phar- maceu- tical Inc.)		
	doxy- cycline; Roche; Periostat		Gingival crevicular fluid collagenase is reported to be inhibited at concentra- tions of 5- 10 microg /ml or 15-

Compound	Trade Name	Reference	Dosage
			30 microM
2S, 5R, 6S-3- aza-4-oxo-10- oxa-5-isobutyl- 2-(N- methylcarbox- amido)- [10]paracyclopha- ne-6-N- hydroxycarboxami- de		WO 97/18207	
hydroxamic acid and amino- carboxylate compounds		WO 96/33176	
N-hydroxamic derivatives of succinamide		WO 96/33166	
Macrocyclic amino carboxylates		J Med Chem 1998 41 11 1749-1751	
	SE-205 (Du Pont Merck Pharm Co.)	Bio-organic Med Chem Lett 1998 8 7 837-842. J Med Chem 1998 41 11 1745 -1748	
macrocyclic matrix metalloprotease-			

Publ. No. WO 00/38717

Compound	Trade Name	Reference	Dosage
8 inhibitors			
Hydroxamic acid and carboxylic acid derivatives		WO 95/22966	
succinamid derivatives		US 5256657	
mercaptosulfide derivatives		WO 95/09833	
sulfoximine and sulfodiimine derivatised peptides		WO 95/09620	
water soluble MMP inhibitors		WO 96/33968	
hydantoin derivatives		EP 06/40594	
Piperazine derivatives		WO 98/27069	
	GI-155704A	J Med Chem 1994 37 5 674. Bioorganic Med Chem Lett 1996 6 16 1905 - 1910	
Cyclic imide derivatives.		EP 05/20573	
3-(mercapto- methyl) hexa-		WO 97/48685	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Compound	Trade Name	Reference	Dosage
hydro-2,5-pyrazinedione derivatives			
beta-mercaptoketone and beta-mercaptoalcohol derivatives		WO 96/40738	
	ilomastat MPI; GM-6001; Galardin	US 5114953. Cancer Res 1994 54 17 4715-4718	eye drops containing ilomastat (800 microg/ml)
Cyclic and heterocyclic N-substituted alpha-iminohydroxamic and carboxylic acids		WO 97/18194	
Aminomethyl-phosphonic and aminomethyl-phosphinic acids derivatives		EP 703239	
3-Mercapto-acetylamino-1,5-substituted-2-oxo-azepan derivatives		WO 98/12211	
2-substituted		WO 94/04531	

Compound	Trade Name	Reference	Dosage
indane-2-mercaptoacetyl-amide tricyclic derivatives			
	Ro-2756 (Roche Holding AG)		
	Ro-26-4325 (Roche Holding AG)		
	Ro-26-5726 (Roche Holding AG)		
	Ro-26-6307 (Roche Holding AG)		
	Ro-31-9790 (Roche Holding AG)	J Am Soc Nephrol 1995 6 3 904. Inflamm Res 1995 44 8 345 -349	mono-arthritis in rat: 100 mg/kg/day
substituted and unsubstituted hydroxamates (specifically N-[D,L-2-isobutyl-		WO 92/09556	

Compound	Trade Name	Reference	Dosage
3-(N'-hydroxy-carbonyl-amido)-propanoyl]tryptophanmethanamide)			
GM6001, N-(2(R)-2 - (hydroxyaminocarboxymethyl)-4-methylpentanoyl)-L-tryptophan methanamide.		WO 95/24921	
Oligonucleotide (c-jun)			
Sulfated polysaccharides		WO 98/11141	
	KB-R7785; KB-R8301; KB-R8845	Life Sci 1997 61 8 795-803	
Fas ligand solubilization inhibitor		WO 97/09066	
gelastatin AB, KRIBB			
	KT5-12 (Kotobuki Seiyaku Co Ltd.)	Faseb J 1998 12 5 A773 (4482)	
2-(N2-[(2R)-2-(2-hydroxyamino-2-oxoethyl)-5-(4-		GB 23/18789	

Publ. No. 00/38717

Compound	Trade Name	Reference	Dosage
methoxyphenoxy)p entanoyl]-L- phenylalanylamin o)ethanesulfonam ide, and carboxylic acid derivatives thereof			
Chromone derivatives		EP 758649	2- Pyrolylthio -chromone in a murine melanoma model produced 37% inhibition at 100 mg/kg
Esculetin derivatives,		EP 719770	
substituted and unsubstituted hydroxyureas and reverse hydroxamates		WO 92/09563	
Synthetic MMP inhibitors (ex. N-(D,L-2- isobutyl-3-(N'- hydroxycarbonyla		WO 94/22309	

Compound	Trade Name	Reference	Dosage
(2S,4R)-2-methyl-4-(phenylamino-carbonylmethyl-aminocarbonyl)-6-(4-propyl-phenyl)hexanoic acid, and carboxylic acid derivatives		WO 97/11936	
Substituted cyclic derivatives		US 5403952	
Thiol sulfonamide metalloprotease inhibitors		WO 98/03166	
Thiol sulfone metalloproteinase inhibitors		WO 98/03164	
formulations containing vanadium compounds and N-acetylcysteine		WO 97/47296	
	NSC-683551; COL-3 (National Cancer Institute)		

Compound	Trade Name	Reference	Dosage
	BB-3644 (Neures Ltd.)		
Arylsulfonamido- substituted hydroxamic acids	CGS- 27023A; CGS-25966	Int Congr Inflamm Res Assoc 1994 7th Abs 73. EP-00606046	600 mg tid (Ph I - colorectal and melanoma patients); 100 mg/kg in food in osteoarthri- tis model rabbits
alpha- Substituted arylsulfonamido hydroxamic acid derivatives		WO 97/22587	
Arylsulfonamido- substituted hydroxamic acids		US 5455258	active at 30 mg/kg in in vivo assay
Arylsulfonamido- substituted hydroxamic acids		WO 96/00214	
2S,3S)-N- hydroxy-5- methyl-2-[2-(2- methoxyethoxy)et hoxymethyl]-3-		WO 98/14424	

Publ. No. 98/00214

Compound	Trade Name	Reference	Dosage
		Lett 1995 5 15 1637 - 1642	
debromohymeni- aldisine and related compounds		WO 96/40147	preferred 1-30 mg/day
amide derivatives of 5-amino-1,3,4- thiadiazolones		WO 96/40745	
3S-(4-(N- hydroxylamino)- 2R- isobutylsuccinyl)amino-1- methoxymethyl- 3,4- dihydrocarbostyr il and derivatives thereof		WO 94/21612	
Carbostyryl derivatives		JP 8325232	
OPB-3206 (Otsuka Pharmaceutical Co, Ltd.)			
Arylsulfonyl hydroxamic acid derivatives		WO 96/33172	
Cyclic sulfone		EP 818442	

Publ. No. WO 00/38717

[illegible]

[illegible]

[illegible]

Compound	Trade Name	Reference	Dosage
amide			
N-hydroxy-phosphinic acid amides		WO 98/08853	
N'-arylsulfonyl derivatives of spirocyclic-N-hydroxycarbox-amides		WO 98/08850	
N'-arylsulfonyl derivatives of thiazepinone and azepinone-N-hydroxycarbox-amides		WO 98/08827	
Substituted piperazine derivatives		WO 98/08825	
N'-arylsulfonyl derivatives of pyrimidine, thiazepine and diazepine-N-hydroxycarbox-amides		WO 98/08823	
Substituted pyrrolidine derivatives		WO 98/08815	
Substituted heterocycles		WO 98/08814	
Substituted 1,3-		WO 09/08822	

PCT/US99/30676

Compound	Trade Name	Reference	Dosage
diheterocyclic derivatives			
substituted 5-amino-1,2,4-thiadiazole-2-thiones		WO 98/25949	
Hydroxamic acid derivatives which inhibit TNF production.		WO 97/24117	
6-methoxy-1,2,3,4-tetrahydro-norharman-1-carboxylic acid		WO 97/37658	
	RS-130830	Arthritis Rheum 1997 40 9 SUPPL. S128	
Aralkyl MMP inhibitors (ex. N-(2R-carboxymethyl-5-(biphen-4-yl)pentanoyl)-L-t-butylglycine-N'-(pyridin-4-yl)carboxamide)		WO 96/16027	
	Ro-32-3555 (Roche Holding		

Compound	Trade Name	Reference	Dosage
	AG)		
	Ro-32-1278 (Roche Holding AG)		
	Ro-32-1541 (Roche Holding AG)		
	Ro-31-3790 (Roche Holding AG)		Arthritic model rats: Protection of cartilage degradation following oral administrat ion; ED50 = 10 mg/kg po
(3R,11S)-N- hydroxy-5- methyl-3-(10- oxo-1,9- diazatricyclo- (11.6.1.014,19)e icosa- 13(20),14(19),15 ,17-tetraen-11- ylcarbamoyl)hexa namide and		WO 95/04735	

Compound	Trade Name	Reference	Dosage
derivatives thereof			
Bridged indoles (Roche Holding AG)		WO 96/23791	
substituted phenylsulfonyl acetamide, propionamide and carboxamide compounds		EP 780386	
5-(4'-biphenyl)-5-[N-(4-nitrophenyl)piperazinyl] barbituric acid		WO 97/23465	
Malonic acid based matrix metalloproteinase inhibitors		EP 716086	
phenyl carboxamide derivatives		WO 95/12603	
Malonic acid based mmp inhibitors (specifically 2-(4-acetylamino-benzoyl)-4-methylpentanoic acid)		EP 716086	

The following individual patent references listed in Table No. 6 below, hereby individually incorporated by reference, describe various MMP inhibitors suitable for use in the present invention described herein, and processes for their manufacture.

10

EP 189784	US 4609667	WO 98/25949	WO 98/25580
JP 10130257	WO 98/17655	WO 98/17645	US 5760027
US 5756545	WO 98/22436	WO 98/16514	WO 98/16506
WO 98/13340	WO 98/16520	WO 98/16503	WO 98/12211
WO 98/11908	WO 98/15525	WO 98/14424	WO 98/09958
WO 98/09957	GB 23/18789	WO 98/09940	WO 98/09934
JP 10045699	WO 98/08853	WO 98/06711	WO 98/05635
WO 98/07742	WO 98/07697	WO 98/03516	WO 98/03166
WO 98/03164	GB 23/17182	WO 98/05353	WO 98/04572
WO 98/04287	WO 98/02578	WO 97/48688	WO 97/48685
WO 97/49679	WO 97/47599	WO 97/43247	WO 97/43240
WO 97/43238	EP 818443	EP 818442	WO 97/45402
WO 97/40031	WO 97/44315	WO 97/38705	US 5679700
WO 97/43245	WO 97/43239	WO 97/43237	JP 09227539
WO 97/42168	US 5686419	WO 97/37974	WO 97/36580
WO 97/25981	WO 97/24117	US 5646316	WO 97/23459
WO 97/22587	EP 780386	DE 19548624	WO 97/19068

[illegible]

WO 93/14199	WO 93/14096	WO 93/13741	WO 93/09090
EP 53/2465	EP 532156	WO 93/00427	WO 92/21360
WO 92/09563	WO 92/09556	EP 48/9579	EP 489577
US 5114953	EP 45/5818	US 5010062	AU 90/53158
WO 97/19075	US 7488460	US 7494796	US 7317407
EP 277428	EP 23/2027	WO 96/15096	WO 97/20824
US 5837696			

The Marimastat used in the therapeutic combinations of the present invention can be prepared in the manner set forth in WO 94/02,447.

5 The Bay-12-9566 used in the therapeutic combinations of the present invention can be prepared in the manner set forth in WO 96/15,096.

The AG-3340 used in the therapeutic combinations of the present invention can be prepared in the manner set
10 forth in WO 97/20,824.

The Metastat used in the therapeutic combinations of the present invention can be prepared in the manner set forth in U.S. Patent No. 5,837,696.

The D-2163 used in the therapeutic combinations of
15 the present invention can be prepared in the manner set forth in WO 97/19,075.

More preferred zinc matrix metalloproteinase inhibitors include those described in the individual U.S. Patent applications, PCT publications and U.S.
20 Patents listed below in Table No. 7, and are hereby individually incorporated by reference.


Table No. 7. More preferred zinc matrix
metalloproteinase inhibitors

U.S. Patent Application Serial Number 97/12,873
U.S. Patent Application Serial Number 97/12,874
U.S. Patent Application Serial Number 98/04,299
U.S. Patent Application Serial Number 98/04,273
U.S. Patent Application Serial Number 98/04,297
U.S. Patent Application Serial Number 98/04,300
U.S. Patent Application Serial Number 60/119,181
WO 94/02447
WO 96/15096
WO 97/20824
WO 97/19075
US 5837696

Even more preferred zinc matrix metalloproteinase inhibitors that may be used in the present invention include:

5

M1)




N-hydroxy-1-(4-methylphenyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;

10

M2)


-73-



1-cyclopropyl-N-hydroxy-4-[[4-[4-(trifluoromethoxy)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;

5


M3)



N-hydroxy-1-(phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;


10

M4)



N-hydroxy-1-(4-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide dihydrochloride;


5 M5)



N-hydroxy-2,3-dimethoxy-6-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]benzamide;

10

M6)




N-hydroxy-1-(4-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide dihydrochloride;

15

M7)


-75-



N-hydroxy-1-(3-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide dihydrochloride;

5


M8)



N-hydroxy-1-(2-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;

10

9)



15


British Biotech BB-2516 (Marimastat), N4-[2,2-dimethyl-1-[(methylamino)carbonyl]propyl]-

09857076 10634

-76-


N1,2 -dihydroxy-3 (2-methylpropyl)-, [2S-
[N4(R*), 2R*, 3S*]]-);

M10)



5 Bayer Ag Bay-12-9566, 4-[(4'-chloro[1,1'-
iphenyl]- 4-yl)oxy]-2-
[(phenylthio)methyl]butanoic acid;

M11)




10

Agouron Pharmaceuticals AG-3340, N-hydroxy-2,2
dimethyl- 4-[[4-(4-pyridinyloxy)phenyl]-
sulfonyl]- 3-thiomorpholinecarboxamide;

15 M12) CollaGenex Pharmaceuticals CMT-3 (Metastat),
6-demethyl-6-deoxy-4-dedimethylaminotetracycline;


M13) Chiroscience D-2163, 2-[1S- ((2R,S)-
20 acetylmercapto-5-phthalimido]pentanoyl-L-
leucyl)amino-3-methylbutyl]imidazole;

M14)




N-hydroxy-4-[[4-(phenylthio)phenyl]sulfonyl]-
1-(2-propynyl)-4-piperidinecarboxamide
monohydrochloride;

M15)




N-hydroxy-1-(2-methoxyethyl)-4-[[4-[4-
(trifluoromethoxy) phenoxy]phenyl]sulfonyl]-4-
piperidinecarboxamide monohydrochloride;

M16)




M17)




1-cyclopropyl-N-hydroxy-4-[[4-[(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;

M18)



4-[[4-(cyclohexylthio)phenyl]sulfonyl]-N-hydroxy-1-(2-propynyl)-4-piperidinecarboxamide monohydrochloride;


M19)



4-[[4-(4-chlorophenoxy)phenyl]sulfonyl]tetrahydro-N-hydroxy-2H-pyran-4-carboxamide;

5


M20)



N-hydroxy-4-[[4-(4-methoxyphenoxy)phenyl]sulfonyl]-1-(2-propynyl)-4-piperidinecarboxamide;

10


M21)



1-cyclopropyl-4-[[4-[(4-fluorophenyl)thio]phenyl]sulfonyl]-N-hydroxy-4-piperidinecarboxamide;

15

M22)




-81-

1-cyclopropyl-N-hydroxy-4-[[4-(phenylthio)phenyl]sulfonyl]-4-piperidinecarboxamide;

5

M23)



10

tetrahydro-N-hydroxy-4-[[4-(4-pyridinylthio)phenyl]sulfonyl]-2H-pyran-4-carboxamide;

M24)



15


tetrahydro-N-hydroxy-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-2H-pyran-4-carboxamide.

20

Still more preferred MMP inhibitors include:


M1)

-82-



N-hydroxy-1-(4-methylphenyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;


M2)



1-cyclopropyl-N-hydroxy-4-[[4-[4-(trifluoromethoxy)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;

5


M3)



N-hydroxy-1-(phenylmethyl)-4-[[4-[4-(trifluoromethoxy)phenoxy]-1-piperidinyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;

10


M4)



15

N-hydroxy-1-(4-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide dihydrochloride;


5 M5)



N-hydroxy-2,3-dimethoxy-6-[[4-[4-(trifluoromethyl)phenoxy]-1-piperidinyl]sulfonyl]benzamide;

10

M6)




N-hydroxy-1-(4-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide dihydrochloride;

15

M7)


-85-



N-hydroxy-1-(3-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide dihydrochloride;

5


M8)



N-hydroxy-1-(2-pyridinylmethyl)-4-[[4-[4-(trifluoromethyl)phenoxy]phenyl]sulfonyl]-4-piperidinecarboxamide monohydrochloride;


10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2



British Biotech BB-2516 (Marimastat), N4-[2,2-dimethyl- 1-[(methylamino)carbonyl]propyl]-N1,2 -dihydroxy-3 (2- methylpropyl)-, [2S-[N4(R*),2R*,3S*]]-);

M10)




10

Bayer Ag Bay-12-9566, 4-[(4'-chloro[1,1'-
iphenyl]- 4-yl)oxy]-2-
[(phenylthio)methyl]butanoic acid;

[illegible]

M11)



Agouron Pharmaceuticals AG-3340, N-hydroxy-
2,2- dimethyl- 4-[[4-(4-pyridinyloxy)phenyl]
sulfonyl]- 3- thiomorpholinecarboxamide;

M12) CollaGenex Pharmaceuticals CMT-3 (Metastat),
6-demethyl-6-deoxy-4-dedimethylaminotetracycline;

M13) Chiroscience D-2163, 2- [1S- ((2R,S)-
acetylmercapto- 5- phthalimido]pentanoyl- L-
leucyl)amino- 3- methylbutyl]imidazole.

Dosage of MMP Inhibitors

Dosage levels of MMP inhibitors on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels of about 1.0 mg to about 1,000 mg. The amount of active ingredient that may be combined with other anticancer agents to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

It is understood, however, that a specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate

of excretion, drug combination, and the severity of the particular disease being treated and form of administration.

Treatment dosages generally may be titrated to
5 optimize safety and efficacy. Typically, dosage-effect relationships from in vitro initially can provide useful guidance on the proper doses for patient administration. Studies in animal models also generally may be used for guidance regarding effective dosages for treatment of
10 cancers in accordance with the present invention. In terms of treatment protocols, it should be appreciated that the dosage to be administered will depend on several factors, including the particular agent that is administered, the route administered, the condition of
15 the particular patient, etc. Generally speaking, one will desire to administer an amount of the compound that is effective to achieve a serum level commensurate with the concentrations found to be effective in vitro. Thus, where an compound is found to demonstrate in vitro
20 activity at, e.g., 10 μ M, one will desire to administer an amount of the drug that is effective to provide about a 10 μ M concentration in vivo. Determination of these parameters are well within the skill of the art.

These considerations, as well as effective
25 formulations and administration procedures are well known in the art and are described in standard textbooks.

Administration Regimen

30 Any effective treatment regimen can be utilized and readily determined and repeated as necessary to effect treatment. In clinical practice, the compositions

containing a MMP inhibitor alone or in combination with other therapeutic agents are administered in specific cycles until a response is obtained.

For patients who initially present without advanced
5 or metastatic cancer, a MMP inhibitor in combination with radiation therapy, is used as a continuous post-treatment therapy in patients at risk for recurrence or metastasis (for example, in adenocarcinoma of the prostate, risk for metastasis is based upon high PSA,
10 high Gleason's score, locally extensive disease, and/or pathological evidence of tumor invasion in the surgical specimen). The goal in these patients is to inhibit the growth of potentially metastatic cells from the primary tumor during surgery and inhibit the growth of tumor
15 cells from undetectable residual primary tumor.

For patients who initially present with advanced or metastatic cancer, a MMP inhibitor in combination with radiation therapy of the present invention is used as a continuous supplement to, or possible replacement for
20 hormonal ablation. The goal in these patients is to slow or prevent tumor cell growth from both the untreated primary tumor and from the existing metastatic lesions.

Illustrations

25 The following discussion highlights some agents in this respect, which are illustrative, not limitative. A wide variety of other effective agents also may be used.

Colorectal Cancer

30 The preferred combination therapy for the treatment of colorectal cancer is surgery, followed by a regimen of one or more chemotherapeutic agents, cycled over a

over a one year time period. In the treatment of colorectal cancer, radiation alone or in combination with surgery and/or chemotherapeutic agents is often used. Preferred chemotherapeutic agents include
5 fluorouracil, and Levamisole. Preferably, fluorouracil and Levamisole are used in combination.

Prostate Cancer

Current therapies for prostate cancer focus upon
10 reducing levels of dihydrotestosterone to decrease or prevent growth of prostate cancer. Radiation alone or in combination with surgery and/or chemotherapeutic agents is often used.

15 Pancreas Cancer

Preferred combinations of therapy for the treatment of non-metastatic adenocarcinoma include the use of preoperative biliary tract decompression (patients presenting with obstructive jaundice); surgical
20 resection, including standard resection, extended or radial resection and distal pancreatectomy (tumors of body and tail); adjuvant radiation; and chemotherapy. For the treatment of metastatic adenocarcinoma, the preferred chemotherapy consists of 5-fluorouracil,
25 followed weekly cisplatin therapy.

Lung Cancer

In many countries including Japan, Europe and America, the number of patients with lung cancer is
30 fairly large and continues to increase year after year and is the most frequent cause of cancer death in both men and women. Although there are many potential causes

for lung cancer, tobacco use, and particularly cigarette smoking, is the most important. Additionally, etiologic factors such as exposure to asbestos, especially in smokers, or radon are contributory factors. Also
5 occupational hazards such as exposure to uranium have been identified as an important factor. Finally, genetic factors have also been identified as another factor that increase the risk of cancer.

Lung cancers can be histologically classified into
10 non-small cell lung cancers (e.g. squamous cell carcinoma(epidermoid), adenocarcinoma, large cell carcinoma (large cell anaplastic), etc.) and small cell lung cancer (oat cell). Non-small cell lung cancer (NSCLC) has different biological properties and
15 responses to chemotherapeutics from those of small cell lung cancer (SCLC). Thus, chemotherapeutic formulas and radiation therapy are different between these two types of lung cancer.

Non-Small Cell Lung Cancer

20 Where the location of the non-small cell lung cancer tumor can be easily excised (stage I and II disease) surgery is the first line of therapy and offers a relatively good chance for a cure. However, in more advanced disease (stage IIIa and greater), where the
25 tumor has extended to tissue beyond the bronchopulmonary lymph nodes, surgery may not lead to complete excision of the tumor. In such cases, the patient's chance for a cure by surgery alone is greatly diminished. Where surgery will not provide complete removal of the NSCLC
30 tumor, other types of therapies must be utilized.

Today radiation therapy is the standard treatment to control unresectable or inoperable NSCLC. Improved results have been seen when radiation therapy has been

combined with chemotherapy, but gains have been modest and the search continues for improved methods of combining modalities.

Radiation therapy is based on the principle that
5 high-dose radiation delivered to a target area will result in the death of reproductive cells in both tumor and normal tissues. The radiation dosage regimen is generally defined in terms of radiation absorbed dose (rad), time and fractionation, and must be carefully
10 defined by the oncologist. The amount of radiation a patient receives will depend on various consideration but the two most important considerations are the location of the tumor in relation to other critical structures or organs of the body, and the extent to
15 which the tumor has spread. A preferred course of treatment for a patient undergoing radiation therapy for NSCLC will be a treatment schedule over a 5 to 6 week period, with a total dose of 50 to 60 Gy administered to the patient in a single daily fraction of 1.8 to 2.0 Gy,
20 5 days a week. A Gy is an abbreviation for Gray and refers to 100 rad of dose.

However, as NSCLC is a systemic disease, and radiation therapy is a local modality, radiation therapy as a single line of therapy is unlikely to provide a
25 cure for NSCLC, at least for those tumors that have metastasized distantly outside the zone of treatment. Thus, the use of radiation therapy with other modality regimens have important beneficial effects for the treatment of NSCLC.

30 Generally, radiation therapy has been combined temporally with chemotherapy to improve the outcome of treatment. There are various terms to describe the temporal relationship of administering radiation therapy

and chemotherapy, and the following examples are the preferred treatment regimens and are generally known by those skilled in the art and are provided for illustration only and are not intended to limit the use of other combinations. "Sequential" radiation therapy and chemotherapy refers to the administration of chemotherapy and radiation therapy separately in time in order to allow the separate administration of either chemotherapy or radiation therapy. "Concomitant" radiation therapy and chemotherapy refers to the administration of chemotherapy and radiation therapy on the same day. Finally, "alternating" radiation therapy and chemotherapy refers to the administration of radiation therapy on the days in which chemotherapy would not have been administered if it was given alone.

It is reported that advanced non-small cell lung cancers do not respond favorably to single-agent chemotherapy and useful therapies for advanced inoperable cancers have been limited. (J. Clin. Oncol. 1992, 10, 829-838).

Japanese Patent Kokai 5-163293 refers to 16-membered-ring macrolide antibiotics as a drug delivery carrier capable of transporting anthracycline-type anticancer drugs into the lungs for the treatment of lung cancers. However, the macrolide antibiotics specified herein are disclosed to be only a drug carrier, and there is no reference to the therapeutic use of macrolides against non-small cell lung cancers.

WO 93/18652 refers to the effectiveness of the specified 16-membered-ring macrolides such as bafilomycin, etc. in treating non-small cell lung cancers, but they have not yet been clinically practicable. Pharmacology, vol. 41, pp. 177-183 (1990)

describes that a long-term use of erythromycin increases productions of interleukins 1, 2 and 4, all of which contribute to host immune responses, but there is no reference to the effect of this drug on non-small cell lung cancers. Tetragenesis, Carcinogenesis, and Mutagenesis, vol. 10, pp. 477-501 (1990) describes that some of antimicrobial drugs can be used as an anticancer agent, but does not refer to their application to non-small cell lung cancers. In addition, interleukins are known to have an antitumor effect, but have not been reported to be effective against non-small cell lung cancers. Any 14 - or 15-membered-ring macrolides have not been reported to be effective against non-small cell lung cancers.

However, several chemotherapeutic agents have been shown to be efficacious against NSCLC. Preferred chemotherapeutic agents against NSCLC include etoposide, carboplatin, methotrexate, 5-fluorouracil, epirubicin, doxorubicin, and cyclophosphamide. The most preferred chemotherapeutic agents active against NSCLC include cisplatin, ifosfamide, mitomycin C, epirubicin, vinblastine, and vindesine.

Other agents that are under investigation for use against NSCLC include: camptothecins, a topoisomerase 1 inhibitor; navelbine (vinorelbine), a microtubule assembly inhibitor; taxol, inhibitor of normal mitotic activity; gemcitabine, a deoxycytidine analogue; fotemustine, a nitrosourea compound; and edatrexate, a antifol.

The overall and complete response rates for NSCLC has been shown to increase with use of combination chemotherapy as compared to single-agent treatment. Haskel, Chest. 1991, 99: 1325; Bakowsk, Cancer Treat.

Rev. 1983; 10:159; Joss, *Cancer Treat. Rev.* 1984, 11: 205.

Small Cell Lung Cancer

Approximately 15 to 20 percent of all cases of lung cancer reported worldwide is small cell lung cancer (SCLC). (Ihde, *Cancer* 1984, 54, 2722). Currently, treatment of SCLC incorporates multi-modal therapy, including chemotherapy, radiation therapy and surgery. Response rates of localized or disseminated SCLC remain high to systemic chemotherapy, however, persistence of the primary tumor and persistence of the tumor in the associated lymph nodes has led to the integration of several therapeutic modalities in the treatment of SCLC.

The most preferred chemotherapeutic agents against SCLC include vincristine, cisplatin, carboplatin, cyclophosphamide, epirubicin (high dose), etoposide (VP-16) I.V., etoposide (VP-16) oral, isofamide, teniposide (VM-26), and doxorubicin. Preferred single-agents chemotherapeutic agents include BCNU (carmustine), vindesine, hexamethylmelamine (altretamine), methotrexate, nitrogen mustard, and CCNU (lomustine). Other chemotherapeutic agents under investigation that have shown activity against SCLC include iroplatin, gemcitabine, lonidamine, and taxol. Single-agent chemotherapeutic agents that have not shown activity against SCLC include mitoguazone, mitomycin C, aclarubicin, diaziquone, bisantrene, cytarabine, idarubicin, mitomxantrone, vinblastine, PCNU and esorubicin.

The poor results reported from single-agent chemotherapy has led to use of combination chemotherapy.

Additionally, radiation therapy in conjunction MMP inhibitors and systemic chemotherapy is contemplated to

be effective at increasing the response rate for SCLC patients. The typical dosage regimen for radiation therapy ranges from 40 to 55 Gy, in 15 to 30 fractions, 3 to 7 times week. The tissue volume to be irradiated
5 is determined by several factors and generally the hilum and subcarinal nodes, and bilateral mediastinal nodes up to the thoracic inlet are treated, as well as the primary tumor up to 1.5 to 2.0 cm of the margins.

Breast Cancer

10 Today, among women in the United States, breast cancer remains the most frequent diagnosed cancer. One in 8 women in the United States at risk of developing breast cancer in their lifetime. Age, family history, diet, and genetic factors have been identified as risk
15 factors for breast cancer. Breast cancer is the second leading cause of death among women.

Different chemotherapeutic agents are known in the art for treating breast cancer. Cytotoxic agents used for treating breast cancer include

20 doxorubicin, cyclophosphamide, methotrexate, 5-fluorouracil, mitomycin C, mitoxantrone, taxol, and epirubicin. (CANCER SURVEYS, Breast Cancer volume 18, Cold Spring Harbor Laboratory Press, 1993).

In the treatment of locally advanced
25 noninflammatory breast cancer, a matrix metalloproteinase inhibitor and radiation therapy can be used to treat the disease in combination with other antiangiogenic agents, or in combination with surgery, or with chemotherapeutic agents. Preferred combinations
30 of chemotherapeutic agents, and surgery that can be used in combination with the radiation therapy and MMP inhibitors include, but are not limited to: 1) doxorubicin, vincristine; 2) cyclophosphamide,

- doxorubicin, 5-flourouracil, vincristine, prednisone; 3) cyclophosphamide, doxorubicin, 5-flourouracil, premarin, tamoxifen; 4) cyclophosphamide, doxorubicin, 5-flourouracil, premarin, tamoxifen, mastectomy; 5) mastectomy, levamisole; 6) mastectomy; and 7) mastectomy, cyclophosphamide, doxorubicin, 5-fluorouracil, tamoxifen, halotestin.

- In the treatment of locally advanced inflammatory breast cancer, MMP inhibitors and radiation therapy can be used to treat the disease in combination with other antiangiogenic agents, or in combination with surgery, or with chemotherapeutic agents. Preferred combinations of chemotherapeutic agents, radiation therapy and surgery that can be used in combination with the MMP inhibitors and radiation include, but or not limited to:
- 1) cyclophosphamide, doxorubicin, 5-fluorouracil; 2) cyclophosphamide, doxorubicin, 5-fluorouracil, mastectomy; 3) 5-fluorouracil, doxorubicin, cyclophosphamide, vincristine, prednisone, mastectomy;
 - 4) 5-fluorouracil, doxorubicin, cyclophosphamide, vincristine, mastectomy; 5) cyclophosphamide, doxorubicin, 5-fluorouracil, vincristine; 6) cyclophosphamide, doxorubicin, 5-fluorouracil, vincristine, mastectomy; 7) doxorubicin, vincristine, methotrexate, followed by vincristine, cyclophosphamide, 5-fluorouracil; 8) doxorubicin, vincristine, cyclophosphamide, methotrexate, 5-fluorouracil, followed by vincristine, cyclophosphamide, 5-fluorouracil; 9) surgery, followed by cyclophosphamide, methotrexate, 5-fluorouracil, prednisone, tamoxifen, followed by cyclophosphamide, methotrexate, 5-fluorouracil, prednisone, tamoxifen, doxorubicin, vincristine, tamoxifen; 10) surgery, followed by cyclophosphamide,

- methotrexate, 5-fluorouracil, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, doxorubicin, vincristine, tamoxifen; 11) surgery, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, followed by cyclophosphamide, methotrexate, 5-fluorouracil, doxorubicin, vincristine, tamoxifen;; 12) surgery, followed by cyclophosphamide, methotrexate, 5-fluorouracil, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, doxorubicin, vincristine; 13) surgery, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, doxorubicin, vincristine, tamoxifen; 14) surgery, followed by cyclophosphamide, methotrexate, 5-fluorouracil, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, doxorubicin, vincristine; 15) surgery, followed by cyclophosphamide, methotrexate, 5-fluorouracil, predinsone, tamoxifen, followed by cyclophosphamide, methotrexate, 5-fluorouracil, doxorubicin, vincristine; 16) 5-fluorouracil, doxorubicin, cyclophosphamide followed by mastectomy, followed by 5-fluorouracil, doxorubicin, cyclophosphamide.

In the treatment of metastatic breast cancer, radiation therapy and MMP inhibitors are used to treat the disease in combination with surgery, or with chemotherapeutic agents. Preferred combinations of chemotherapeutic agents, and surgery that can be used in combination with the radiation therapy and MMP inhibitors include, but are not limited to: 1) cyclophosphamide, methotrexate, 5-fluorouracil; 2)

- cyclophosphamide, adriamycin, 5-fluorouracil; 3) cyclophosphamide, methotrexate, 5-fluorouracil, vincristine, prednisone; 4) adriamycin, vincristine; 5) thiotepa, adriamycin, vinblastine; 6) mitomycin, 5 vinblastine; 7) cisplatin, etoposide.

Bladder Cancer

The classification of bladder cancer is divided into three main classes: 1) superficial disease, 2) muscle-invasive disease; and 3) metastatic disease.

- 10 Currently, transurethral resection (TUR), or segmental resection, account for first line therapy of superficial bladder cancer, i.e., disease confined to the mucosa or the lamina propria. However, intravesical therapies are necessary, for example, for the treatment
15 of high-grade tumors, carcinoma in situ, incomplete resections, recurrences, and multifocal papillary. Recurrence rates range from up to 30 to 80 percent, depending on stage of cancer.

- Therapies that are currently used as intravesical
20 therapies include chemotherapy, immunotherapy, bacille Calmette-Guerin (BCG) and photodynamic therapy. The main objective of intravesical therapy is twofold: to prevent recurrence in high-risk patients and to treat disease that cannot be resected. The use of
25 intravesical therapies must be balanced with its potentially toxic side effects. Additionally, BCG requires an unimpaired immune system to induce an antitumor effect. Chemotherapeutic agents that are known to be inactive against superficial bladder cancer
30 include Cisplatin, actinomycin D, 5-fluorouracil, bleomycin, and cyclophosphamide methotrxate.

In the treatment of superficial bladder cancer, MMP inhibitors and radiation therapy are used to treat the

disease in combination with surgery (TUR), and intravesical therapies.

Preferred combinations of chemotherapeutic agents are selected from the group consisting of thiotepa (30
5 to 60 mg/day), mitomycin C (20 to 60 mg/day), and doxorubicin (20 to 80 mg/day).

The preferred intravesicle immunotherapeutic agent that may be used in the present invention is BCG. The preferred daily dose ranges from 60 to 120 mg, depending
10 on the strain of the live attenuated tuberculosis organism used.

The preferred photodynamic therapeutic agent that may be used with the present invention is Photofrin I, a photosensitizing agent, administered intravenously. It
15 is taken up by the low-density lipoprotein receptors of the tumor cells and is activated by exposure to visible light. Additionally, neodymium YAG laser activation generates large amounts of cytotoxic free radicals and singlet oxygen.

20 In the treatment of muscle-invasive bladder cancer, radiation therapy and MMP inhibitors can be used to treat the disease in combination with other antiangiogenic agents, or in combination with surgery (TUR), intravesical chemotherapy, and radical cystectomy
25 with pelvic lymph node dissection.

The preferred radiation dose is between 5,000 to 7,000 cGY in fractions of 180 to 200 cGY to the tumor. Additionally, 3,500 to 4,700 cGY total dose is administered to the normal bladder and pelvic contents
30 in a four-field technique. Radiation therapy should be considered only if the patient is not a surgical candidate, but may be considered as preoperative therapy.

The preferred combination of chemotherapeutic agents that can be used in combination with radiation therapy and the MMP inhibitors is cisplatin, methotrexate, vinblastine.

5 Currently no curative therapy exists for metastatic bladder cancer. The present invention contemplates an effective treatment of bladder cancer leading to improved tumor inhibition or regression, as compared to current therapies.

10 In the treatment of metastatic bladder cancer, a combination of radiation therapy and MMP inhibitors can be used to treat the disease in combination with surgery, or with chemotherapeutic agents.

Preferred combinations of chemotherapeutic agents
15 include, but are not limited to: 1) cisplatin and methotrexate; 2) doxorubicin, vinblastine, cyclophosphamide, and 5-fluorouracil; 3) vinblastine, doxorubicin, cisplatin, methotrexate; 4) vinblastine, cisplatin, methotrexate; 5) cyclophosphamide,
20 doxorubicin, cisplatin; 6) 5-fluorouracil, cisplatin.

Head and Neck Cancers

Head and neck cancer accounts for approximately 2% of new cancer cases in the United States. Common intracranial neoplasms include glioma, meningioma,
25 neurinoma, and adenoma. Preferred combinations that can be used along with a combination of radiation therapy and an integrin antagonist for the treatment of malignant glioma include: 1) BCNU (carmustine);
2) methyl CCNU (lomustine); 3) medrol; 4) procarbazine;
30 5) BCNU, medrol; 6) misonidazole, BCNU;
7) streptozotocin; 8) BCNU, procarbazine; 9) BCNU, hydroxyurea, procarbazine, VM-26; 10) BNCU, 5-fluorouracil; 11) methyl CCNU, dacarbazine;

12) misonidazole, BCNU; and 13) PCNU. The preferred dose of radiation therapy is about 5,500 to about 6,000 cGY. Preferred radiosensitizers include misonidazole, intra-arterial Budr and intravenous iododeoxyuridine
5 (IUDR).

Biological Evaluation

Solitary tumors are generated in the right hind legs of mice by the injection of 3×10^5 viable NFSA tumor cells. Treatment with a MMP inhibitor (6 mg/kg
10 body weight) or vehicle (0.05% Tween 20 and 0.95% polyethylene glycol) given in the drinking water is started when tumors are approximately 6 mm in diameter and the treatment is continued for 10 consecutive days. Water bottles are changed every 3 days. Tumor
15 irradiation is performed 3-8 days after initiation of the treatment with a MMP inhibitor. The end points of the treatment are tumor growth delay (days) and TCD₅₀ (tumor control dose 50, defined as the radiation dose yielding local tumor cure in 50% of irradiated mice 120
20 days after irradiation). To obtain tumor growth curves, three mutually orthogonal diameters of tumors are measured daily with a vernier caliper, and the mean values are calculated.

Local tumor irradiation with single γ -ray doses of
25 30, 40, or 50 Gy is given when these tumors reach 8 mm in diameter. Irradiation to the tumor is delivered from a dual-source ¹³⁷Cs irradiator at a dose rate of 6.31 Gy/minute. During irradiation, unanesthetized mice are immobilized on a jig and the tumor is centered in a
30 circular radiation field 3 cm in diameter. Regression and regrowth of tumors are followed at 1-3 day intervals until the tumor diameter reaches approximately 14 mm.